

D-A186 111

OFFICE OF NAVAL RESEARCH

Contract N00014-86-0043

TECHNICAL REPORT No. 52

MTC FILE COPY

Coherent States for the Damped Harmonic Oscillator

Þу

k. H. Yeon, C. I. Um and Thomas F. George

Prepared for Publication

. .

Phishat Review A

Cepartments of Chemistry and Physics State University of New York at Buffalo Buffalo, New York (4.5)

THE CONTRACT OF THE

Tell duling in whole on in lant is permitted for any purpose of the inited States Government.

This document has been approved for public release and sale; its distribution is unlimited.



	REPORT DOCUME	NTATION PAGE	E			
IN REPORT SECURITY CLASSIFICATION Unclassified	16. RESTRICTIVE MARKINGS					
28 SECURITY CLASSIFICATION AUTHORITY		Approved for public release; distribution				
26 DECLASSIFICATION DOWNGRADING SCHEDULE		unlimited				
4 PERFORMING ORGANIZATION REPORT NUM	BER(S)	5. MONITORING ORGANIZATION REPORT NUMBER(\$)				
UBUFFALO/DC/87/TR-52						
Depts. Chemistry & Physics State University of New York	5b OFFICE SYMBOL (If applicable)	78 NAME OF MONIT	TORING ORGAN	ZATION		
Fronczak Hall, Amherst Campus Buffalo, New York 14260		76 ADDRESS (City, State and AIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217				
Be NAME OF FUNDING SPONSORING ORGANIZATION	Bb OFFICE SYMBOL ilf applicable:	9 PROCUREMENT I			MBEA	
Office of Naval Research	<u> </u>	Contract N00014-86-K-0043		043		
Lhemistry Program 800 N. Quincy Street Arlington, Virginia 22217		PROGRAM ELEMENT NO	PROJECT NO	TASK NO.	WORK UNIT	
Coherent States for t	he Damped Harmor	nic Oscillator				
12 PERSONAL AUTHORIS) K. H. Yeon.	C. I. Um and Tho	omas F. George	· · · · · · · · · · · · · · · · · · ·			
134 TYPE OF REPORT 136 TIME C		14 DATE OF REPOR		15 PAGE CO	TAUC	
Prepared for publication in Physical Review A						
FIELD GROUP SUB GR	DAMPED HARMONIC COHERENT STATES	OSCILLATOR	CALDIRO	A-KANAI HAM DISSIPATIVE	ILTONIAN "	
19 ASSTRACT Continue on reverse if necessary and						
Using the Caldirola-Kanai	Hamilitonian for	tne damped n	armonic osc	chilator, ex	act	
coherent states are cons	tructed. These	new coherent	states sat	isfy the pro	perties	
which coherent states should generally have.						
			,			
·						
		•				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT		21 ABSTRACT SECURITY CLASSIFICATION				
UNCLASSIFIED/UNLIMITED & SAME AS APT. & DTIC USERS		Unclassified				
224. NAME OF RESPONSIBLE INDIVIDUAL		22b TELEPHONE NUMBER 22c OFFICE SYMBOL (Include Area Code)		30L		
Dr. David L. Nelson		(202) 696-44	10			

Coherent States for the Damped Harmonic Oscillator

K. H. Yeon and C. I. Um*
Department of Physics, College of Science, Korea University
Seoul 136, Korea

Thomas F. George
Departments of Chemistry and Physics
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260 USA

Using the Caldirola-Kanai Hamiltonian for the damped harmonic oscillator, exact coherent states are constructed. These new coherent states satisfy the properties which coherent states should generally have.



PACS Nos. 02.90,, 03.65 W, 05.30-d, 42.50. Bs

* To whom correspondence should be addressed

Access	ion For	
MIIS	GRADI	3
DTIC 1		
Unanno		
Justii	lication	
	ibution/	Codes
Dist	Avail a	
DIAC	3001	
10/		
111	1 1	
(1	

Since the coherent states for the harmonic oscillator have been constructed first by Schrödinger, 1) they have been widely used to describe many fields of physics. 2)-5) Recently Nieto and Simmons have constructed coherent states for particles in general potentials () and have applied their formalism to confining one dimensional systems: 7) harmonic oscillator with centripetal barrier and the symmetric Pöschl-Teller potential, and also to nonconfining one dimensional systems 8) with the symmetric Rosen-Morse potential and the Morse potential. For the time-dependent systems Lewis and Riesenfeld have investigated the harmonic oscillator with timedependent frequency $\omega(t)$. Hartley and Ray $\frac{10}{2}$ has obtained exact coherent states for this time-dependent harmonic oscillator on the basis of Lewis and Riesenfeld theory. Hartley-Ray results satisfy most, but not all, of the properties of the coherent states. In the case of a quantum mechanical model of a damped forced harmonic oscillator, Dodonov and Man'koll) have introduced the Caldirola-Kanai Hamiltonian 12) with an external force term and constructed integrals of motion of this Hamiltonian, eigenstates and coherent states. The main flaw of the Dodonov-Man'ko result is its uncertainty relation $\Delta p \cdot \Delta x \ge e^{-\gamma t} h/2$ in which the uncertainty vanishes as $t \rightarrow \infty$. This contradiction is critically reviewed by Greenberger 13), and Cervero and Villarroel. (14) Greenberger introduced the variable mass: m = m_ne^{7t} and removed the violation of uncertainty.

In this paper we construct exact coherent states for the damped harmonic oscillator described by the Caldirola-Kanai Hamiltonian

$$\mathcal{X} = e^{-\gamma t} \frac{p^2}{2m} + e^{\gamma t} \frac{1}{2} m \omega_0^2 x^2.$$
 (1)

We first define creation operator a and annihilation operator a and using these operators we will derive the representations of coherent states and investigate whether our coherent states satisfy the following properties of coherent states: (1) They are eigenstates of annihilation operator.

(2) They are created from the vacuum or the ground states by an unitary operator. (3) They represent the minimum uncertainty states.

They are not orthogonal but complete and normalized.

In the preceeding paper 15 (hereafter referred to as paper I) we have developed the quantum theory of the damped driven harmonic oscillator with the Caldirola-Kanai Hamiltonian with an external driving force f(t) by path integral method. In paper I, setting f(t) = 0, the Hamiltonian is reduced to Eq. (1) and all other results become those corresponding to the Hamiltonian (Eq.(1)), and the Lagrangian, mechanical energy and propagator are given by

$$f = e^{\gamma t} (\frac{1}{2} m x^2 - \frac{1}{2} m \omega_0^2 x^2), \qquad (2)$$

$$E = e^{-27t} \frac{p^2}{2m} + \frac{1}{2} m \omega_0^2 x^2, \qquad (3)$$

$$K(x,t; x_0, 0) = \left(\frac{\frac{7}{2}t}{2\pi i \hbar \sin \omega t}\right)^{\frac{1}{2}} exp\left[\frac{im}{4\hbar} \left\{\gamma(x_0^2 - e^{\gamma t} x^2)\right\} + \frac{2\omega}{\sin \omega t} \left((x^2 e^{\gamma t} + x_0^2) \cos \omega t - 2e^{\frac{\gamma}{2}t} xx_0\right)\right\},$$
(4)

with $\omega = (\omega_0^2 - \gamma^2/4)^{\frac{1}{2}}$. Here, the energy expression in Eq. (3) is not equal to the Hamiltonian itself. With the help of Eq. (4) and the wave function of simple harmonic oscillator, we obtain the wave function of the

damped harmonic oscillator:

$$\psi_{n}(x,t) = \frac{N}{\sqrt{2^{n} n!}} H_{n}(Dx) \exp\{-i(n + \frac{1}{2})\cot^{-1}(\frac{7}{2\omega} + \cot \omega t) - Ax^{2}\},$$
(5)

where

$$N = \left(\frac{m\omega}{\pi h}\right)^{\frac{1}{4}} \frac{e^{-\frac{1}{4}7t}}{\frac{1}{5(t)\sin^{\frac{1}{2}}\omega t}},$$

$$S^{2}(t) = \frac{7^{2}}{4\omega^{\frac{1}{2}}} + \frac{7}{\omega}\cot\omega t + \csc^{2}\omega t,$$

$$A(t) = \frac{m\omega}{2h} e^{7t} \left\{ \frac{1}{\frac{1}{5(t)^{2}\sin^{2}\omega t}} + i\left(\frac{7}{2\omega} - \cot\omega t + \frac{\frac{7}{2\omega} + \cot\omega t}{\frac{7}{5(t)^{2}\sin^{2}\omega t}}\right) \right\},$$

$$D(t) = \left(\frac{m\omega}{h}\right)^{\frac{1}{4}} \frac{e^{\frac{1}{2}7t}}{\frac{7}{5(t)\sin\omega t}}.$$
(6)

The quantum mechanical expectation values of mechanical energy E(Eq. (3)) take the form

$$< E>_{mn} = -\frac{h^2}{2m} e^{-2\gamma t} < \frac{\partial^2}{\partial x^2}>_{mn} + \frac{1}{2} m \omega_0^2 < x^2>_{mn}.$$
 (7)

The evaluation of Eq. (7) gives the non-zero matrix elements which occur only in the principal diagonal and the two second off-diagonals:

$$_{n+2,n} = [(n+2)(n+1)]^{\frac{1}{2}} \theta(t),$$
 (8)
 $_{nn} = \frac{1}{2} (n + \frac{1}{2})\hbar\omega e^{-\gamma t} \left[\frac{\omega_0^2}{\omega^2} \zeta(t)^2 \sin^2\omega t + \frac{1}{\zeta(t)^2 \sin^2\omega t}\right],$

where

$$\theta(t) = \frac{1}{4} \hbar \omega e^{-\gamma t} \exp \left\{ \left[2i \cot^{-1} \left(\frac{\gamma}{2\omega} + \cot \omega t \right) \right] \left[\int_{0}^{\infty} (t)^{2} \sin^{2} \omega t \right] - \frac{1}{\left[\int_{0}^{\infty} (t)^{2} \sin^{2} \omega t \right]} + \frac{1}{\left[\int_{0}^{\infty} (t)^{2} \sin^{2} \omega t \right]} \left[\left(\frac{\gamma}{2\omega} - \cot \omega t \right) \int_{0}^{\infty} (t)^{2} \sin^{2} \omega t \right] + \frac{\gamma}{2\omega} + \cot \omega t \right\} - 2i \left\{ \left(\frac{\gamma}{2\omega} - \cot \omega t \right) \int_{0}^{\infty} (t)^{2} \sin^{2} \omega t \right\} + \cot \omega t \right\} \right\}.$$

$$(10)$$

Taking the complex conjugate and changing n into (n-2) in Eq. (8) we can easily obtain the energy expectation value in (n-2, n) state.

In a similar way to that used to obtain Eq. (7) we can obtain the uncertainty relations in the various states:

$$[(\Delta p)(\Delta x)]_{n+2,n} = \frac{\hbar}{2} \{(n+2)(n+1)\}^{\frac{1}{2}} \beta(t),$$
 (11)

$$[(\Delta p)(\Delta x)]_{n+1,n} = \frac{h}{2}(n+1)\beta(t),$$
 (12)

$$\left[\Delta p \cdot \Delta x\right]_{nn} = \left(n + \frac{1}{2}\right) h\beta(t), \qquad (13)$$

and

$$\beta(t) = \left[1 + \left\{ \left(\frac{1}{8} \left(\frac{\gamma}{\omega}\right)^3 + \left(\frac{\gamma}{\omega}\right)\right) \sin^2 \omega t + \frac{1}{8} \left(\frac{\gamma}{\omega}\right)^2 \sin 2\omega t \right\}^2 \right]^{\frac{1}{2}}.$$
(14)

Chainging n into (n-1) and (n-2) respectively in Eqs. (12) and (11) we can obtain the uncertainty relations in (n-1, n) and (n-2, n) states.

Before we construct annihilation operator a and creation operator a,

These
we give the properties of the coherent states. The coherent states can be
defined by the eigenstates of the nonhermitian operator a,

$$a|\alpha\rangle = \alpha|\alpha\rangle. \tag{15}$$

Using the completeness relation for the number representations, we can expand $|\alpha>as$

$$|\alpha > = e^{-\frac{1}{2}|\alpha|^{2}} \sum_{n=0}^{\infty} \frac{\alpha^{n}}{\sqrt{n!}} |n >$$

$$= e^{-\frac{1}{2}|\alpha|^{2}} e^{\alpha a^{*}} |0 > , \qquad (16)$$

where $|0\rangle$ is the vacuum or ground state and is independent of n. The calculation of $<\beta |\alpha\rangle$ in Eq. (16) gives

$$\langle \beta | \alpha \rangle = e^{-\frac{1}{2}(|\alpha|^2 + |\beta|^2) + \alpha \beta^*}$$
 (17)

Since Eq. (17) has nonzero value for $\alpha \neq \beta$, the states are not orthogonal, but when $|\alpha - \beta|^2 \to \infty$, the states become orthogonal.

The eigenvalue α of coherent states are complex number u + iv, and thus the completeness relation of coherent states is written as

$$\int |\alpha| > \langle \alpha| \frac{d^2 \alpha}{\pi} = 1 , \qquad (18)$$

where 1 is the identity operator and $d^2\alpha$ is given by d(Reu)d(Imv).

To define a^+ and a for the damped harmonic oscillator we make use of Eq. (5) for $\langle x \rangle_{mn}$ and $\langle p \rangle_{mn}$:

$$< x >_{mn} = \int_{-\infty}^{\infty} \psi_{m}^{*}(x) x \psi_{n}(x) dx$$

$$= \frac{1}{2} (n+1)^{\frac{1}{2}} (ReA)^{-\frac{1}{2}} exp\{i \cot^{-1}(\frac{\gamma}{2\omega} + \cot \omega t)\} \delta_{m,n+1}$$

$$+ \frac{1}{2} n^{\frac{1}{2}} (ReA)^{-\frac{1}{2}} exp\{-i \cot^{-1}(\frac{\gamma}{2\omega} + \cot \omega t)\} \delta_{m,n-1}$$

$$= (n + \frac{1}{2})^{\frac{1}{2}} \mu(t) \delta_{m,n+1} + n^{\frac{1}{2}} \mu(t)^{*} \delta_{m,n-1} ,$$

$$_{mn} = \int_{-\infty}^{\infty} \psi_{m}^{*}(x) \frac{h}{i} \frac{\partial}{\partial x} \psi_{n}(x) dx$$

$$= i \hbar \sqrt{2} (n+1)^{\frac{1}{2}} \frac{A}{D} exp\{i \cot^{-1}(\frac{\gamma}{2\omega} + \cot \omega t)\} \delta_{m,n+1}$$

$$+ i \hbar \sqrt{2} n^{\frac{1}{2}} (\frac{A}{D} - D) exp\{-i \cot^{-1}(\frac{\gamma}{2\omega} + \cot \omega t)\} \delta_{m,n-1}$$

$$= (n + \frac{1}{2})^{\frac{1}{2}} \eta(t) \delta_{m,n+1} + n^{\frac{1}{2}} \eta^{*}(t) \delta_{m,n-1},$$

$$(20)$$

where

$$\mu(t) = \frac{1}{2} (ReA)^{-\frac{1}{2}} \exp[i \cot^{-1}(\frac{\gamma}{2\omega} + \cot \omega t)],$$
 (21)

$$\eta(t) = \sqrt{2} i \hbar \frac{A}{D} \exp[i \cot^{-1}(\frac{\gamma}{2\omega} + \cot \omega t)], \qquad (22)$$

and we have the relation

$$\eta \mu^* - \eta^* \mu = 2 i \operatorname{Im} \left\{ \frac{1}{2} \left(\operatorname{ReA} \right)^{+\frac{1}{2}} \sqrt{2} i \hbar \frac{A}{D} \right\} = i \hbar.$$
 (23)

Therefore, we define annihilation operator a and creation operator a for the damped harmonic oscillator as follows

$$a = \frac{1}{i\hbar} (\eta x - \mu p), \qquad (24)$$

$$a^* = \frac{1}{i\hbar} (\mu^* p - \eta^* x),$$
 (25)

when the expressions of x and p by a and a are

$$x = \mu^* a + \mu a^*,$$
 (26)

$$p = \eta^* a + \eta a^*.$$
 (27)

Since η is not equal to μ in Eqs. (21) - (22), we can easily conform that a and a^+ are not hermitian operators, but the following relations are preserved:

$$[x,p] = ih$$
 (28)

$$[a,a^{\dagger}] = 1.$$
 (29)

Now we evaluate the transformation function $< x | \alpha >$ from coherent states to the coordinate representation | x > . From Eqs. (15) and (24)

we have

$$(\eta x' - \mu \frac{h}{i} \frac{\partial}{\partial x'}) < x' | \alpha > = i h \alpha < x' | \alpha > , \qquad (30)$$

For convenience we change the variable x' into x and solve this differential to whom equation, and we get

$$< x | \alpha > = N \exp \{ \frac{1}{\mu} \alpha x - (2 i \hbar \mu)^{-1} \eta x^{2} \},$$
 (31)

where N is the integral constant. Taking N to satisfy Ey. (18), we obtain the eigenvectors of operator a given in the coordinate representation |x>:

$$< x | \alpha > = (2 \pi \mu \mu^{*})^{-\frac{1}{2}} exp \{ -\frac{1}{2ih} \frac{\eta}{\mu} x^{2} + \frac{\alpha}{\mu} x - \frac{1}{2} |\alpha|^{2} - \frac{1}{2} \frac{\mu^{*}}{\mu} \alpha^{2} \}$$
 (32)

Next we show that a coherent state represents a minimum uncertainty state. With the help of the relations between a, a^+ , x and p we evaluate the expectation values of x, p, x^2 and p^2 in state $|\alpha>$:

$$\langle x \rangle = \langle \alpha | \mu^{+} a + \mu a^{+} | \alpha \rangle = \mu^{+} \alpha + \mu \alpha^{+},$$

$$\langle p \rangle = \langle \alpha | \eta^{+} a + \eta a^{+} | \alpha \rangle = \eta^{+} \alpha + \eta \alpha^{+},$$

$$\langle x^{2} \rangle = \mu^{+2} \alpha^{2} + \mu \mu^{+} (1 + 2\alpha \alpha^{+}) + \mu^{2} \alpha^{+2},$$

$$\langle p^{2} \rangle = \eta^{+2} \alpha^{2} + \eta \eta^{+} (1 + 2\alpha \alpha^{+}) + \eta^{2} \alpha^{+2}.$$

$$(33)$$

From Eq. (33) we have

$$(\Delta x)^{3} = \langle x^{3} \rangle - \langle x \rangle^{2} = \mu \mu^{*}, \qquad (34)$$

$$(\Delta p)^2 = \langle p^2 \rangle - \langle p \rangle^2 = \eta \eta^*,$$
 (35)

and thus the uncertainty relation becomes

$$(\Delta x)(\Delta p) = \{|\eta|^2 |\mu|^2\}^{\frac{1}{2}} = \frac{h}{2} \beta(t).$$
 (36)

Eq. (36) is the minimum uncertainty corresponding to Eq. (13) in (0,0) state.

All of the formulas, we have derived, are reduced to those of simple harmonic oscillator when $\tau = 0$. The propagator (Eq. (4)) has very similar form to those of Cheng 15) and others, 17) but the wave function (Eq. (5)) is of new form.

We should note that the same classical equation of motion can be obtained from many different actions, and thus one may have many different propagators corresponding to the actions.— Therefore it is very important to get the correct propagator. The mechanical energy (Eq. (3)) is not Hawke identical to the Hamiltonian operator (Eq. (1)). Therefore we assume that this Hamiltonian represents the quantum mechanical dissipative system.

Fig. 1 and 2 illustrate the decay of the energy expectation value and Atthough the uncertainty relation as a function of γ/ω in (n, n) state. As though we have shown only the principal diagonal element, i.e., $\langle E \rangle_{nn}$ of the energy expectation values, there are four off-diagonals adjacent to the principal diagonal, which are involved in the exponential decaying term

 $e^{-\gamma t}$. $\langle E \rangle_{nn}$ approaches when the constant value as $\frac{\gamma}{\omega} \to 0$. The uncertainty for (n, n) state with period π (Eq. (13)) is reduced to that of the harmonic oscillator at 0° and 180° .

From all of the above we conclude that the coherent states for the damped harmonic oscillator, with the Caldirola-Kanai Himiltonian we have constructed, satisfy the properties of coherent states (1)-(4).

Acknowledgements

We are


I am grateful to Professor S. W. Kim for helpful comments. This research was supported partly by Korea University and by the office of Naval Research, the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009, and the National Science Foundation under Grant CHE-8620274.


Figure Caption

- Fig. 1. Energy expectation value for (n, n) state as a function of ω t at the various value of γ/ω . As γ/ω tends to zero, energy approaches the constant values.
- Fig. 2. Uncertainty relation for (n, n) state versus ω t at we various values of γ/ω .

References

- 1. E. Schrödinger, Naturwiss. 14, 166 (1926).
- 2. Z.E. Zimmerman and A.H. Silver, Phys. Rev. 167, 418 (1968).
- R.J. Glauber, Phys. Rev. 130, 2529 (1963): ibid 131, 2766 (1963);
 W.H. Louisell, Quantum Statistical Properties of Radiation (John Wiley and Sons, New York, 1973);
 M. Sargent III, M.O. Scully and W.E. Lamb, Laser Physics (Addision-Wesley, New York, 1974).
- J.S. Langer, Phys. Rev. <u>167</u>, 183 (1968); F.W. Cummings and J.R. Johnston, Phys. Rev. 151, 105 (1966).
- 5. J.C. Botke, D.J. Scalapino, and R.L. Sugar, Phys. Rev. D9, 813 (1974).
- 6. M.M. Nieto and L.M. Simmons, Jr., Phys. Rev. D20, 1321 (1979).
- 7. M.M. Nieto and L.M. Simmons, Jr., Phys. Rev. D20, 1342 (1979).
- 8. M.M. Nieto and L.M. Simmons, Jr., Phys. Rev. D20, 1342 (1979).
- 9. H.R. Lewis, Jr., and W.B. Riesenfeld, J. Math. Phys. 10, 1458 (1969).
- J.G. Hartley and J.R. Ray, Phys. Rev. D25, 382 (1982).
- 11. V.V. Dodonov and V.I. Manko, Phys. Rev. A20, 550 (1979).
- P. Coldirola, Nuovo Cimento, 18, 393 (1941); ibid. B77, 2475 (1983);
 E. Kanai, Prog. Theor. Phys. 3, 440 (1948).
- 13. D.M. Greenberger, J. Math. Phys. 20, 762 (1979).
- 14. J.J. Cervero and J. Villarroel, J. Phys. A: Math. Gen. 17, 2963 (1984).
- C.I. Um, K.H. Yeon and W.H. Kahng, J. Phys. A: Math. Gen. <u>20</u>, 611 (1987).
 K.H. Yeon, C.I. Um and W.H. Kahng, New Physics <u>25</u>, 107 (1985);
 ibid. 25, 117 (1985).
- 16. B.K. Cheng, J. Phys. A: Math. Gen. 17, 2475 (1985).
- 17. D.C. Khandekar and S.V. Lawanda, J. Math. Phys. 16, 384 (1975).





01/1113/86/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies	•	No. Copies
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

Dr. J. E. Jensen Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
Chemistry Depa
27709
George Washing

Dr. M. Grunze Laboratory for Surface Science and Technology University of Maine Orono, Maine 04469

Dr. J. Butler Naval Research Laboratory Code 6115 Washington D.C. 20375-5000

Dr. L. Interante Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Irvin Heard Chemistry and Physics Department Lincoln University Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde Department of Chemistry Kansas State University Manhattan, Kansas 66506 Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookesville, Tennessee 38501

Dr. D. Dilella Chemistry Department George Washington University Washington D.C. 20052

Dr. R. Reeves Chemistry Department Renssaeler Polytechnic Institute Troy, New York 12181

Dr. Steven M. George Stanford University Department of Chemistry Stanford, CA 94305

Dr. Mark Johnson Yale University Department of Chemistry New Haven, CT 06511-8118

Dr. W. Knauer Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Dr. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720

Dr. J. Murday Naval Research Laboratory Code 6170 Washington, D.C. 20375-5000

Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Theodore E. Madey Surface Chemistry Section Department of Commerce National Bureau of Standards Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne Chemistry Department Northwestern University Evanston, Illinois 60637

Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712

Dr. D. E. Harrison Department of Physics Naval Postgraduate School Monterey, California 93940 Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green Quantum Surface Dynamics Branch Code 3817 Naval Weapons Center China Lake, California 93555

Dr. A. Wold Department of Chemistry Brown University Providence, Rhode Island 02912

Dr. S. L. Bernasek Department of Chemistry Princeton University Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

Dr. F. Carter Code 6170 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. Richard Colton Code 6170 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. Dan Pierce National Bureau of Standards Optical Physics Division Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer Materials Characterization Lab. General Electric Company Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee R301 Naval Surface Weapons Center White Oak Silver Spring, Maryland 20910

Or. Paul Schoen Code 6190 Naval Research Laboratory Washington, D.C. 20375-5000 Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene Code 5230 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda University of Pittsburg Chemistry Building Pittsburg, PA 15260

Dr. E. A. Irene Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514

Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974

Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5MH UNITED KINGDOM

Dr. H. Tachikawa Chemistry Department Jackson State University Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

Dr. R. G. Wallis Department of Physics University of California Irvine, California 92664

Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052

Dr. J. C. Hemminger Chemistry Department University of California Irvine, California 92717

Dr. T. F. George Chemistry Department University of Rochester Rochester, New York 14627

Dr. G. Rubloff IBM Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598

Dr. Horia Metiu Chemistry Department University of California Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser Department of Chemistry University of Richmond Richmond, Virginia 23173

Dr. R. W. Plummer Department of Physics University of Pennsylvania Philadelphia, Pennsylvania 19104

Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann Department of Chemistry Cornell University Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, NewYork 12181

Dr. G.H. Morrison Department of Chemistry Cornell University Ithaca, New York 14853 END DATE FILMED DEC. 1987